MAX

DOCUMENTED-INPUT ESTIMATES (% w/v, elemental)

N 9.0% **P 0.9%** **K approx. 2.6%** **S 8.3-8.8%**

NET CONTENTS

1,000 L

LIQUID NPKS FERTILISER WITH A BIOLOGICAL BASE | PASTURE, CROPPING & ORCHARDS

Calculated from supplied ingredient data; excludes unquantified inputs and full batch variation. Not a guaranteed finished-product analysis.
Nutrient content, pH and liquid density: pending latest laboratory results; retesting by **31 October 2026.**

PURPOSE & INGREDIENTS

Liquid nutrition for periods of active pasture or crop nutrient demand. Contains fish hydrolysate, molasses, charged biochar, seaweed extract, soluble mineral fertiliser and water.

BIOCHARGE RECOMMENDATIONS

Apply as part of a planned liquid nutrition programme. Match rate and frequency to crop type, growth stage, soil condition, testing and the existing fertiliser programme.

Use suitable application equipment and follow normal good fertiliser practice. **Agitate thoroughly before use and maintain agitation during application.**

Contact Biocharge to confirm water volume, compatibility, equipment suitability and the current application advice. Follow the label and wear appropriate PPE.

Grazing / harvest interval: obtain MAX-specific advice from Biocharge before use.

HAZARD STATEMENTS

H302: Harmful if swallowed.
H319: Causes serious eye irritation.
H433: Harmful to terrestrial vertebrates.

HANDLING PRECAUTIONS

WARNING
Acute oral toxicity Category 4; eye irritation Category 2.
HSNO: **HSR002571**. Read the hazard statements and precautions before use.

Keep away from children. Avoid swallowing, skin/eye contact and spray mist. **Wear chemical-resistant gloves, splash goggles and protective clothing.** Use suitable respiratory protection where mist cannot be controlled. Wash after use; do not eat, drink or smoke during handling.

LIVESTOCK & ENVIRONMENT

Contains fish-derived material; not for human or animal consumption. Keep animals away from stored product and spills.

Excessive copper or selenium can be toxic to livestock. Obtain veterinary or suitably qualified advice before giving supplements or veterinary medicines containing these elements to animals grazing treated pasture.

Prevent overspray, spills and runoff entering waterways. Avoid excessive application or localised nutrient accumulation.

FIRST AID

Eyes: Rinse gently for at least 15 minutes. Remove lenses if easy. Obtain medical assessment.
Skin: Rinse and remove contaminated clothing. Seek advice if irritation or a rash develops.
Inhaled: Fresh air; seek help if unwell.
Swallowed: Clear the mouth. Do not induce vomiting. Call the Poisons Centre. No drinks unless advised; nothing by mouth if unconscious.

Poisons & Medical Exposure Advice:
National Poisons Centre
0800 764 766 (0800 POISON)

STORAGE, SPILLS & DISPOSAL

Store cool, dry and well ventilated, out of sunlight. Keep labelled, upright and secure, away from food/feed, strong acids, alkalis and oxidisers.

Vent the lid every 30 days. Inspect more often if the storage area gets hot. If the tank is bulging, contact Biocharge before opening.

Contain spills and collect residues for appropriate disposal. Keep runoff out of waterways. Do not reuse containers for food, feed or drinking water.

BATCH NUMBER

DATE OF MANUFACTURE

BEST BEFORE: 24 MONTHS FROM DATE OF MANUFACTURE

Refer to the current MAX safety data sheet.
Also available in 200 L drums.

Manufactured by Biocharge Ltd
121 Kana Street, Mataura, New Zealand
021 819 372 | info@biocharge.co.nz

biocharge.co.nz