

WARNING
Keep out of reach of children
ECOTOXIC

Rainbow & Brown

DICAMBA 500 SL

A selective herbicide for the control of certain broadleaf weeds in conservation tillage programs and in cereals, maize, some forage brassicas, pasture and waste areas. (Controls atrazine resistant fathen in maize)

ACTIVE INGREDIENT:

Contains 500 g/l dicamba as the Dimethylamine Salt in the form of a soluble concentrate.

Contains

5
litres



DATE OF
MANUFACTURE:

BATCH No:

EMERGENCY RESPONSE: Call 0800 CHEMCALL, or 0800 243 622 (24 hours)

DICAMBA 500 SL

READ THE LABEL AND LEAFLET COMPLETELY BEFORE MIXING OR APPLYING

HSNO CLASSIFICATIONS: Skin irritation Category 2, Eye irritation Category 2, Specific target organ toxicity (repeated exposure) Category 2, Hazardous to aquatic environment, acute and chronic Category 1, Hazardous to soil organisms, to terrestrial vertebrates and to terrestrial invertebrates. **WARNING:** Causes serious skin or eye irritation. May cause liver damage from repeated ingestion of high doses.

ECOTOXIC : Very toxic to aquatic life with long-lasting effects. Must not be applied directly onto or into water. Very toxic to the soil environment. Toxic to terrestrial vertebrates and terrestrial invertebrates. This product must be under the control of a suitably qualified handler when mixed, loaded or applied. Take all reasonable steps to ensure that the product does not cause any significant adverse effects to the environment beyond the application area. Do not apply product to plants if – (1) Bees are foraging; or (2) The plants are in flower or part flower, and are likely to be visited by non-target invertebrate pollinators (including bees).

PRECAUTIONS: PERSONAL PROTECTION: Do not eat, drink or smoke while using. Wear protective clothing such as impervious gloves, waterproof hat, coat and trousers when using. Remove protective clothing and wash hands and face thoroughly before meals and after work. Avoid contact with the eyes. Wash personal protective equipment after use.

STORAGE: Keep out of reach of children. Store in original container tightly closed and in a secure dry, cool, well ventilated area, away from food stuffs. Store in accordance with the current version of NZS 8409 Management of Agrichemicals. Store containing 100 L of **RAINBOW & BROWN DICAMBA 500 SL** require secondary containment and signage, and more than 100 L requires an emergency response plan.

SPILLS AND DISPOSAL: When dealing with spills wear personal protective clothing and equipment as described in the PERSONAL PROTECTION section. Collect up spilled material, into labelled drums that can be sealed for safe disposal. If spray mixture is spilled, contain and absorb with suitable inert absorbent material and collect in drums as above. Deal with all spillages immediately. If contamination of drains, streams, watercourses, etc. is unavoidable, contact the local regulatory authority. **Product Disposal:** Dispose of this product only by using according to the label or at an approved landfill. Avoid contamination of any water supply with product or empty container. **Container Disposal:** Ensure container is completely empty. Triple rinse container and add residue to mixture. Recycle containers if possible, otherwise bury in an approved land fill. Burn if circumstances, especially wind direction, permit. Otherwise bury in a suitable land fill.

FIRST AID: If swallowed DO NOT induce vomiting. Rinse mouth. For advice, contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately. If on skin, remove contaminated clothing and wash affected areas with plenty of water. DO NOT SCRUB THE SKIN. If in eyes, flush with running water for at least 15 minutes. Remove contact lenses if present after the first 5 minutes and keep rinsing. If eye irritation persists, get medical advice. If inhaled, remove to fresh air and keep at rest until recovered. Get medical advice if person feels unwell.

Safety Data Sheet: Additional information is provided in the safety data sheet.

MIXING / COMPATIBILITY: Partially fill the spray tank and add the measured quantity of **Rainbow & Brown Dicamba 500 SL**. Agitate while filling and spraying. **Rainbow & Brown Dicamba 500 SL** is compatible with most commonly used herbicides: check the label of the companion product before mixing.

CLEANING: After use flush the spray equipment thoroughly. If the sprayer is to be used on sensitive crops ensure it is thoroughly decontaminated.

GENERAL INFORMATION: **Rainbow & Brown Dicamba 500 SL** is active against many broadleaf weeds and is particularly useful in assisting the activity of phenoxy herbicides such as MCPA 750, 2,4-D ester, 2,4-D amine. **Rainbow & Brown Dicamba 500 SL** can be mixed with glyphosate where control of white clover is desired. Best control is achieved when **Rainbow & Brown Dicamba 500 SL** is applied to seedling weeds (3-leaf stage, or rosettes of less than 5 cm diameter).

• Do not spray if rain is expected within six hours. • Do not spray if weeds are wet with dew or rain, or are under stress from drought, extreme cold or waterlogging. • Do not spray outside recommended crop growth stage do not spray Swedes, forage rape or hybrid brassicas. • **Rainbow & Brown Dicamba 500 SL** is applied to seedling weeds (3-leaf stage, or rosettes of less than 5 cm diameter).

PLANT BACK PERIOD: Do not plant legume crops (peas, beans, lucerne, lupins, clover) within 28 days after spraying **Rainbow & Brown Dicamba 500 SL**. Grass, cereal and brassica crops do not require any waiting period.

WEEDS CONTROLLED OR SUPPRESSED: Annuals: Mouse-ear chickweed, Black medic, Black nightshade, Cape weed, chamomiles, chickweed, cleavers, annual clovers, cornbind, cut-leaved geranium, dove's foot, fathen (N.B. not the dicamba resistant biotype), fireweed, hemp nettle, hedge mustard, mallows, Onehunga weed, nipplewort, prostrate amaranth, redroot (amaranth), shepherd's purse, stinking mayweed, spiny emex, spurrey, thorn apple, vetches, willow weed (redshank), wireweed.

Biennials, Perennials: Alligator weed, Californian thistle, cats ear, dandelion, docks, fennel, bindweed, goat's rue, greater bindweed, hawkbit, hemlock, inkweed, leafy spurge, mallows, pearlwort, pennyroyal, perennial clovers, perennial sow thistle, ragwort, selfheal, sheep's sorrel, wild carrot, wild onion, yarrow.

Weeds not controlled: **Rainbow & Brown Dicamba 500 SL** does not control fumitory, plantains, star weed, stonecrop or speedwells.

Withholding period: It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residues Levels for Agricultural Compounds. Do not graze treated areas or feed treated crops or threshings to livestock for two weeks after application.

Registered pursuant to the ACVM Act 1997, No P9919

See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to HSNO Act 1996, HSR000442

See www.epa.govt.nz for approval conditions

WARRANTY: This formulation is warranted to contain the specified active ingredient within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors beyond the manufacturer's control. The manufacturer declines all liability for damage resulting from improper storage or use of the product.

Registrant:

Rainbow & Brown Ltd
68a Tallyho Street, Rotorua 3015
P O Box 100 49, Mail Centre, Rotorua 3046
Freecall 0508 299 299. www.RainbowBrown.co.nz



Rainbow & Brown